



PEDIATRIC BRADYCARDIA WITH PULSES 'SYMPTOMATIC'

BASIC EMT / ADVANCED EMT

1. Chest compressions if:
 - Infant heart rate is less than 60 bpm.

PARAMEDIC

2. **Epinephrine**
 - **1:10,000** 0.01mg/kg (0.1 ml/kg) IV / IO or **1:1000** 0.1 mg/kg (0.1 ml/kg) ET tube.
 - Repeat every 4 minutes until blood pressure is adequate for age.
3. **Atropine**
 - 0.02 mg/kg,
 - a. maximum 0.5 mg
 - May repeat once if increased vagal tone or primary AV block.
4. Consider cardiac pacing.

NORMAL PEDIATRIC VITALS

AGE	RESPIRATORY RATE	HEART RATE	BLOOD PRESSURE
Infant (1 to 12 mo)	30 to 60	100 to 160	greater than 60 mm Hg or strong central pulse
Toddler (1 to 3 yr)	24 to 40	90 to 150	greater than 70 mm Hg or strong central pulse
Preschooler (4 to 5 yr)	22 to 34	80 to 140	greater than 75 mm Hg
School-age (6 to 12 yr)	18 to 30	70 to 120	greater than 80 mm Hg
Adolescent (13 to 18 yr)	12 to 16	60 to 100	greater than 90 mm Hg